

# INTRAVENEOUS IMMUNE GLOBULINS

BRAND NAME	Gammagard S/D		IGIVnex & GAMUNEX®	Gammagard Liquid	Privigen®	Panzyga®
WEBLINK	<a href="http://www.baxalta.ca/products_en.html">http://www.baxalta.ca/products_en.html</a>		<a href="http://www.grifols.com">www.grifols.com</a>	<a href="http://www.baxalta.ca/products_en.html">http://www.baxalta.ca/products_en.html</a>	<a href="http://www.cslbehring.ca">www.cslbehring.ca</a>	<a href="http://www.octapharma.ca">www.octapharma.ca</a>
SIZES	5g		5g, 10g, 20g <b>Only 20g for IGIVnex</b>	2.5g, 5g, 10g, 20g, 30g	2.5g, 5g, 10g, 20g, 40g	10g, 20g, 30g
INDICATIONS	PID, CLL, ITP		PID, SID, ITP, CIDP, GBS	PID, SID, ITP, MMN	PID, SID, ITP, CIDP	PID, SID, ITP
CONCENTRATION	5% or 10% upon reconstitution		10%	10%	10%	10%
IgA CONTENT	≤ 2.2 µg/mL in 5% solution		Not more than 0.084mg/mL	≤ 0.14 mg/mL	No more than 25 mcg/mL	≤0.3 mg/mL
SUGAR CONTENT	20 mg/mL dextrose in 5%		Not indicated in product monograph	None	Not indicated in product monograph	Not indicated in product monograph
STORAGE TEMPERATURE	Not to exceed +25°C		+2°C to +8°C	+2 to +8°C	+2°C to +25°C	+2°C to +8 °C
INITIAL Recommended INFUSION RATE	<u>5% &amp; 10% Solution</u> 0.5 mL/kg of body weight/hr		0.01 to 0.02 mL/kg/ min (1 to 2 mg/kg/min) for the first 30 mins	0.5 mL/kg of BW/hr for 30 mins BW=body weight	0.5 mg/kg/min (0.3 mL/kg/hr)	1 mg/kg/ min for the first 30 mins
MAXIMUM Recommended INFUSION RATE	<u>5% Solution</u> 4 mL/kg/hr	<u>10% Solution</u> 8 mL/kg/hr	0.14 mL/kg/min (14 mg/kg/min)	8 mL/kg of BW/hr  <u>For MMN</u> 5.4 mL/kg of body weight/hr	12 mg/kg/min (7.2 mL/kg/hr)	8 mg/kg/min (in new patients)  <u>For PID &amp; SID</u> 12 or 14 mg/kg/min (in experienced patients)

2018-03-20

This Table is a summary and will not tell you everything about IG products. **DETAILED information can be found in the PRODUCT MONOGRAPH or PRODUCT INSERT.** *CONTENT is subject to change.*